Oncology Clinical Pathways Salivary Gland Cancer

March 2024 - V2.2024







Table of Contents

alivary Gland Cancer Presumptive Conditions	3
alivary Gland T1-2, N0, M0.	4
alivary Gland T3-4a, N0.	5
alivary Gland T1-4 Node Positive	
alivary Gland T4b	
alivary Gland Recurrent or Metastatic Disease	
lolecular Testing Table	9





<u>Salivary Gland Cancer – Presumptive Conditions</u>

VA automatically presumes that certain disabilities were caused by military service. This is because of the unique circumstances of a specific Veteran's military service. If a presumed condition is diagnosed in a Veteran within a certain group, they can be awarded disability compensation.

Atomic Veterans Exposed to Ionizing Radiation

Salivary gland cancer

Gulf War and Post 9/11 Veterans

If the patient served any amount of time in Afghanistan, Djibouti, Syria, or Uzbekistan during the Persian Gulf War, from Sept. 19, 2001, to the present or the *Southwest Asia theater of operations from Aug. 2, 1990, to the present, specific conditions include:

- Salivary gland-type tumors of the lung
- Salivary gland-type tumors of the trachea

For more information, please visit <u>U.S. Department of Veterans Affairs - Presumptive Disability Benefits (va.gov)</u>








^{*} The Southwest Asia theater of operations refers to Iraq, Kuwait, Saudi Arabia, the neutral zone between Iraq and Saudi Arabia, Bahrain, Qatar, the United Arab Emirates, Oman, the Gulf of Aden, the Gulf of Oman, the Persian Gulf, the Arabian Sea, the Red Sea, and the airspace above these locations.

Salivary Gland Cancer – T1-2, N0, M0



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.


- ^a **Adverse Features** include positive margins, neural invasion, lymphovascular invasion
- ^b Resection surgical operations may be staged if needed dependent on pathology of primary tumor
- ^c High or Intermediate Grade includes adenoid cystic
- d Surveillance consider resection if low morbidity
- ^e Adenoid Cystic consider long-term surveillance > 5 years







<u>Salivary Gland Cancer – T3-4a, N0</u>



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email ClinicalTrialsNavigation@va.gov.


- ^a Resection surgical operations may be staged if needed dependent on pathology of primary tumor
- ^b **Neck Dissection** consider avoiding if subsequent radiation to the primary site includes a large measure of the nodal basin; radiation fields would need to be increased to provide therapeutic coverage of the neck
- ^c Adverse Features Include intermediate or high grade, positive margins, neural invasion, multiple lymph nodes, advanced T stage (T3-4), extranodal extension, adenoid cystic, or lymphovascular invasion







<u>Salivary Gland Cancer – T1-4 Node Positive</u>



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.


^a **Adverse Features Include** intermediate or high grade, positive margins, neural invasion, multiple lymph nodes, advanced T stage (T3-4), extranodal extension, adenoid cystic, or lymphovascular invasion







Salivary Gland Cancer – T4b



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.


^a **Adverse Features Include** intermediate or high grade, positive margins, neural invasion, multiple lymph nodes, advanced T stage (T3-4), extranodal extension, adenoid cystic, or lymphovascular invasion







<u>Salivary Gland Cancer – Recurrent or Metastatic Disease</u>



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email Clinical trial, email Clinical trial, email ClinicalTrialsNavigation@va.gov.

- ^a Molecular Testing to include HER2 by IHC, NTRK gene fusion, MSI High, and Androgen Receptor; NGS is preferred
- b HER2 includes IHC, considered positive if 3+; if 2+ perform FISH testing
- ^c Evidence Base is limited; limited data from phase 2 clinical trials; abiraterone is an acceptable alternative
- d Leuprolide prescribe for both men and women; in women, either pre or post menopause
- ^e Cisplatin, Doxorubicin, and Cyclophosphamide evaluate cardiovascular risk factors with baseline LVEF







<u>Salivary Gland Cancer – Molecular Testing Table</u>

Eligibility	Test Category	Test Type	Recommended Vendors	NPOP Coverage	Specimen Type
Recurrent or Metastatic Salivary Gland Cancer		DNA and RNA-based comprehensive genomic profiling (CGP)	· ·		Tumor Tissue**, Blood
	IHC	AR HER2 (with reflex to FISH if 2+)	Local VA or locally contracted vendor		Tumor Tissue
	FISH	HER2 FISH if IHC is 2+	Local VA or locally contracted vendor	No	Tumor Tissue

^{*} Somatic NGS should include TMB and MSI and adequately cover NTRK fusions







^{**} Tissue testing strongly preferred because it is the only method for RNA based testing. Liquid testing is suboptimal but acceptable only if adequate tissue cannot be obtained