

Oncology Clinical Pathways Bladder Cancer (Urothelial Carcinoma Only)

July 2025 – V3.2025



Choose **VA**



SHOULDER to SHOULDER
Every Step of the Way

VA



U.S. Department
of Veterans Affairs

Table of Contents

Presumptive Conditions	3
Clinical Presentation and Evaluation	4
Non-Muscle Invasive Surgical Evaluation	5
Non-Muscle Invasive Low Risk	6
Non-Muscle Invasive Intermediate Risk	7
Non-Muscle Invasive High Risk	8
Positive Urine Cytology Only	9
Urothelial Carcinoma of Prostatic Urethra	10
Upper GU Tract Carcinoma	11
Stage II, IIIA, IIIB Muscle Invasive	12
Stage IVA	13
Stage IVB First Line	14
Stage IVB Second Line	15
Post Cystectomy Follow Up	16
Muscle Invasive Post Bladder Preservation Follow Up	17
Molecular Testing Table	18



Bladder Cancer – Presumptive Conditions

VA automatically presumes that certain disabilities were caused by military service. This is because of the unique circumstances of a specific Veteran's military service. If a presumed condition is diagnosed in a Veteran within a certain group, they can be awarded disability compensation.

Vietnam Veterans – Agent Orange Exposure or Specified Locations

- Bladder cancer

Atomic Veterans – Exposure to Ionizing Radiation

- Cancer of the urinary tract

Gulf War and Post 9/11 Veterans


If the patient served on or after Sept. 11, 2001, in Afghanistan, Djibouti, Egypt, Jordan, Lebanon, Syria, Uzbekistan, or Yemen or if the patient served in the *Southwest Asia theater of operations, or Somalia, on or after Aug. 2, 1990, specific conditions include:

- Urinary bladder, ureter, and related genitourinary cancers

*The Southwest Asia theater of operations refers to Iraq, Kuwait, Saudi Arabia, the neutral zone between Iraq and Saudi Arabia, Bahrain, Qatar, the United Arab Emirates, Oman, the Gulf of Aden, the Gulf of Oman, the Persian Gulf, the Arabian Sea, the Red Sea, and the airspace above these locations.

For more information, please visit [U.S. Department of Veterans Affairs - Presumptive Disability Benefits \(va.gov\)](#); [VA News - Presumptive for Service Connection Jan 08 2025](#); [eCFR :: 38 CFR 3.320a -- Presumptive service connection for bladder, ureter, and related genitourinary cancers.](#)

Bladder Cancer – Clinical Presentation and Evaluation



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **CT Urogram** in patients unable to receive IV contrast, order alternative upper tract imaging

^b **Exposure** includes Agent Orange, burn pits, and other occupational/environmental toxins

^c **Cytology** order if results would change clinical management

^d **Family History** family or personal malignancy history, suspicion for Lynch syndrome, or age under 60 years


^e **TURBT with EUA** include blue-light cystoscopy if clinically appropriate

^f **Intravesical Gemcitabine** for known or presumed low grade

TURBT Transurethral Resection of Bladder Tumor

EUA Exam Under Anesthesia

Bladder Cancer – Non-Muscle Invasive Surgical Evaluation




Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Variant Histology** includes micropapillary, nested, plasmacytoid, neuroendocrine, sarcomatoid, squamous or glandular predominant

TURBT Transurethral Resection of Bladder Tumor

American Urological Association Non-Muscle Invasive Risk Stratification		
Low Risk	Intermediate Risk	High Risk
<ul style="list-style-type: none"> Papillary urothelial neoplasm of low malignant potential <p style="text-align: center;">Or</p> <ul style="list-style-type: none"> Low grade urothelial carcinoma <ul style="list-style-type: none"> Ta and ≤3 cm and Solitary 	<ul style="list-style-type: none"> Low grade urothelial carcinoma <ul style="list-style-type: none"> T1 or >3 cm or Multifocal or Recurrence within 1 year <p style="text-align: center;">Or</p> <ul style="list-style-type: none"> High grade urothelial carcinoma <ul style="list-style-type: none"> Ta and <3 cm and Solitary 	<ul style="list-style-type: none"> High grade urothelial carcinoma <ul style="list-style-type: none"> CIS or T1 or >3 cm or Multifocal <p style="text-align: center;">Or</p> <ul style="list-style-type: none"> Very high risk features (any) <ul style="list-style-type: none"> BCG unresponsive Variant histologies ^a Lymphovascular invasion Prostatic urethral involvement


Bladder Cancer – Non-Muscle Invasive Low Risk



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

TURBT Transurethral Resection of Bladder Tumor

Bladder Cancer – Non-Muscle Invasive Intermediate Risk




Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Intravesical Therapy** gemcitabine should begin within 3-4 weeks of TURBT; BCG or gemcitabine maintenance for one year

^b **Cystoscopy Post TURBT Schedule** at Year 1: at 3, 6, and 12 months; Year 2: every 6 months; Years 3 and later: annually

TURBT Transurethral Resection of Bladder Tumor

Bladder Cancer – Non-Muscle Invasive High Risk



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **BCG Naïve** BCG non-exposed or greater than one year since last BCG

^b **Very High Risk Features** include variant histologies, lymphovascular invasion, or prostatic urethral invasion

^c **BCG Induction** only one repeat induction BCG course

^d **Cystoscopy and Cytology Post TURBT** surveillance schedule: years 1-2: every 3 months; years 3-4: every 6 months; years ≥ 5 : annually

^e **BCG Maintenance** 3 week instillations at 3, 6, 12, 18, 24, 30, and 36 months (3 years) after start of induction BCG

^f **Adequate BCG** defined as ≥ 5 induction doses and ≥ 2 maintenance doses

^g **BCG Unresponsive** defined as persistent high-grade disease or recurrence within 6 months of receiving at least 2 courses of intravesical BCG (at least 5 of 6 induction and at least 2 of 3 maintenance doses of BCG)


^h **Pembrolizumab** option for treatment of patients with BCG-unresponsive, high-risk NMIBC with T1s tumors who are ineligible for or have elected not to undergo cystectomy

ⁱ **Nadofaragene Firadenovec** all criteria must be met: BCG unresponsive, non-muscle invasive bladder cancer, Carcinoma in situ (CIS) with or without papillary tumors (Ta or T1 high-grade tumors)

BCG Bacillus Calmette Guerin

TURBT Transurethral Resection of Bladder Tumor

Bladder Cancer – Positive Urine Cytology Only




Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Cytology** review clinical history with cytopathologist

^b **Cystoscopy** use enhanced technology if available

Bladder Cancer – Urothelial Carcinoma of Prostatic Urethra



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Urothelial Carcinoma of Prostatic Urethra** solely isolated prostatic urethra involvement


^b **Needle Biopsy** if DRE or PSA is abnormal

^c **Ta, Tis, T1** based on transurethral resection; confirm subepithelial involvement

^d **If Only Prior Pelvic Metastatic Disease**, reimaging with PET to ensure no metastatic disease prior to proceeding with surgery

TURP Transurethral Resection of the Prostate
BCG Bacillus Calmette-Guerin


Bladder Cancer – Upper GU Tract Carcinoma



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

- ^a **Cystoscopy and Ureteroscopy** may include selective washing ± single-dose intravesical gemcitabine
- ^b **Family History** family or personal malignancy history, suspicion for Lynch syndrome; age under 60 years
- ^c **Mitomycin Gel Instillation** use for ureteral tumors is off-label
- ^d **Extent of Disease** staging to include recent chest and abdominal cross-sectional imaging
- ^e **Consider Neoadjuvant Gemcitabine and Cisplatin** for select high grade patients; consider tumor board discussion

Bladder Cancer – Stage II, IIIA, IIIB Muscle Invasive



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Imaging** perform bone scan if clinically indicated


^b **Candidate for Cystectomy** patients with clinical node positive disease should have resolution of adenopathy post chemo to become eligible for cystectomy

^c **Bladder Preservation** avoid bladder preservation in patients with hydronephrosis and extensive or multifocal carcinoma in situ

^d **SDM Durvalumab** shared decision making is critical at the time of consideration of durvalumab; in clinical trials, the control arm was chemotherapy alone and contained no information regarding any subsequent therapies in the control arm; thus, it is unclear if survival benefit is from neoadjuvant therapy alone, adjuvant therapy, or both components; physician should discuss these uncertainties with patients

MVAC Methotrexate, Vinblastine, Doxorubicin, Cisplatin

Bladder Cancer – Stage IVA



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Imaging** perform bone scan if clinically indicated


^b **Eligible for EV** exclude patients with preexisting peripheral neuropathy \geq Grade 2, baseline ocular disorders, uncontrolled diabetes at baseline

^c **SDM Nivolumab** shared decision making is critical at the time of consideration of nivolumab; in the CheckMate 901 trial, only 40% of patients within the control arm received a checkpoint inhibitor as subsequent therapy; providers will need to assess cisplatin candidacy prior to using this regimen

^d **Interim Assessment** includes EUA, cystoscopy, and CT chest/abdomen/pelvis

CGP Comprehensive Genomic Profiling
EV Enfortumab Vedotin
EUA Exam Under Anesthesia

Bladder Cancer – Stage IVB First Line



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Imaging** perform bone scan if clinically indicated, imaging of central nervous system (CNS) as clinically indicated

^b **Candidate for EV** exclude patients with preexisting peripheral neuropathy \geq Grade 2, baseline ocular disorders, uncontrolled diabetes at baseline


CGP Comprehensive Genomic Profiling

EV Enfortumab Vedotin

Criteria for Use

Enfortumab Vedotin: exclude patients with preexisting neuropathy \geq Grade 2, baseline ocular disorders, or uncontrolled diabetes at baseline

Bladder Cancer – Stage IVB Second Line



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **HER2 IHC3+** that progressed on previous therapy with no satisfactory alternative

Criteria for Use

Erdafitinib: exclude patients with retinal/corneal abnormality at baseline or serum phosphate greater than upper limits of normal at baseline; perform ophthalmological exams at baseline and then monthly for the first 4 months of therapy, then every 3 months thereafter

Bladder Cancer – Post Cystectomy Follow Up

	Year 1	Year 2	Year 3	Year 4	Year 5	Year 5-10	Year >10
Post Cystectomy Follow Up	Imaging NMIBC	CT urogram at 3 & 12 mo	Annual CT urogram			Annual renal ultrasound	As clinically indicated
	Imaging MIBC	CT chest and CT urogram every 3-6 mo		Annual CT chest/abdomen/pelvis		Annual renal ultrasound	As clinically indicated
	Blood Tests	CMP & CBC every 6 mo	Annual CMP and B ₁₂ levels			Annual B ₁₂ levels	
	Urine Tests	Urine cytology every 6-12 mo; consider urethral wash every 6-12 mo		Urine cytology as clinically indicated Urethral wash cytology as clinically indicated ^a			

Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

^a **Urethral Wash** recommended for prostatic urethral involvement (papillary or CIS) on TURBT or cystectomy pathology

Bladder Cancer – Muscle Invasive Post Bladder Preservation Follow Up

Muscle Invasive
Post Bladder Preservation Follow Up

	Year 1	Year 2	Year 3	Year 4	Year 5	Year 5-10	Year >10
Cystoscopy	Every 3 mo		Every 6 mo		Annually		As clinically indicated
Imaging	CT chest and CT urogram every 3-6 mo		Annual CT chest/abdomen/pelvis			As clinically indicated	
Blood Tests	CMP & CBC every 6 mo	Annual CMP					
Urine Tests	Urine cytology every 6-12 mo		Urine cytology as clinically indicated				

Bladder Cancer – Molecular Testing Table

Eligibility	Test Category	Test Type	Recommended Vendors	NPOP Coverage	Specimen Type
Stage IVA Muscle Invasive Urothelial Carcinoma/Bladder Cancer, Predominantly Urothelial	Somatic NGS	CGP using both DNA and RNA based methodology	Tempus Foundation Medicine	Yes Yes	Tumor Tissue*, Blood
	IHC	PD-L1 expression by IHC using 22C3 antibody (pembrolizumab), SP142 antibody (atezolizumab), 28-8 pharmDx antibody (nivolumab), SP263 antibody (durvalumab)	Tempus Foundation Medicine	Yes (When ordered with CGP) Yes (When ordered with CGP)	Tumor Tissue
Stage IVB Metastatic Urothelial Carcinoma/Bladder Cancer	Somatic NGS	CGP using both DNA and RNA based methodology	Tempus Foundation Medicine	Yes Yes	Tumor Tissue*, Blood
	IHC	PD-L1 expression by IHC using 22C3 antibody (pembrolizumab), SP142 antibody (atezolizumab), 28-8 pharmDx antibody (nivolumab), SP263 antibody (durvalumab)	Tempus Foundation Medicine	Yes (When ordered with CGP) Yes (When ordered with CGP)	Tumor Tissue
	IHC	HER2	Foundation Medicine	Yes (When ordered with CGP)	Tumor Tissue

* Tissue preferred, but liquid acceptable if tissue insufficient